
Citation:

Tee, JC and Coopoo, Y and Lambert, MI (2015) Rugby union movement patterns: The impact of fatigue and substitute players. In: 20th Annual Congress of the European College of Sports Medicine, 24 June 2015 - 27 June 2015, Malmo, Sweden.

Link to Leeds Beckett Repository record:

<https://eprints.leedsbeckett.ac.uk/id/eprint/4474/>

Document Version:

Conference or Workshop Item (Presentation)

Creative Commons: Attribution 4.0

The aim of the Leeds Beckett Repository is to provide open access to our research, as required by funder policies and permitted by publishers and copyright law.

The Leeds Beckett repository holds a wide range of publications, each of which has been checked for copyright and the relevant embargo period has been applied by the Research Services team.

We operate on a standard take-down policy. If you are the author or publisher of an output and you would like it removed from the repository, please [contact us](#) and we will investigate on a case-by-case basis.

Each thesis in the repository has been cleared where necessary by the author for third party copyright. If you would like a thesis to be removed from the repository or believe there is an issue with copyright, please contact us on openaccess@leedsbeckett.ac.uk and we will investigate on a case-by-case basis.

Rugby union movement patterns: The impact of fatigue and substitute players

Jason C. Tee^a, Mike I. Lambert^b and Yoga
Coopoo^a

^a Department of Sport and Movement Studies, University of Johannesburg
^b Division of Exercise Science and Sports Medicine, University of Cape Town

Email: jasonctee@gmail.com




Fatigue in team sports

Fatigue = ↓ in total and high-intensity running distance

(Waldron and Highton, 2014, Sports Med 44:12)

Distribution of energy resources



Pacing schema

Macro-pacing (pre-match)

- hydration, fuel availability, motivation, temperature, opposition, whole-game/substitute

Meso-pacing (half time)

- homeostatic disturbance, opposition, scoreline

Micro-pacing (continuous)

- homeostatic disturbance, opposition, scoreline

Edwards and Noakes, 2009, Sports Med 39:1

Professional Rugby Union

Rugby union is characterised by short-duration, high-intensity efforts during which players collide, often while running at full speed; interspersed by longer low-intensity periods of standing, walking and jogging.

(Austin *et al.*, 2011, J Sci Med Sport 14:3)



Diversity of Physical Requirements



The game demands differ for players in different positions.

(Deutsch *et al.*, 2007, J Sport Sci 25:4)

Research Aim

Understand the nature of fatigue in professional rugby union

- What is the influence of match period and position on movement patterns?
- What is the influence of substitutes on movement patterns?

Methods – Global Positioning System (GPS)

Variables measured

- Playing time
- Relative distance ($\text{m} \cdot \text{min}^{-1}$) in speed zones

Speed bands		
Walking	$0-2\text{m} \cdot \text{s}^{-1}$	Low intensity running $0-4\text{m} \cdot \text{s}^{-1}$
Jogging	$2-4\text{m} \cdot \text{s}^{-1}$	
Striding	$4-6\text{m} \cdot \text{s}^{-1}$	High intensity running $>4\text{m} \cdot \text{s}^{-1}$
Sprinting	$>6\text{m} \cdot \text{s}^{-1}$	

- Sprint ($>6\text{m} \cdot \text{s}^{-1}$) frequency
- Acceleration ($>2.75\text{m} \cdot \text{s}^{-2}$) frequency
- Accelerometer
 - Total impacts $>5\text{G} \cdot \text{min}^{-1}$
 - High-intensity impacts $>8\text{G} \cdot \text{min}^{-1}$

SPI Pro GPS unit

(GPSports, Canberra)
mass = 76g;
size = 87 x 48 x 20 mm
5Hz GPS Tracking
100Hz Tri-axial
Accelerometer



Methods



Player characteristics

Age 25.5 ± 2.4 years

Body mass 101.5 ± 12.2 kg

Stature 1.86 ± 0.07 m


- Whole game players – start game and complete >35 min in 2nd half
- Substitute players – 2nd half replacements

Statistics

- Factorial ANOVA
- Paired and independent sample t-tests
- Cohen's effect size




Results – Effect of half on total and high-intensity distance




* indicates significant difference from 1st half. S, M, L and VL indicate effect sizes small (0.2-0.5), medium (0.5-0.8), large (0.8-1.2) and very large (>1.2) respectively.

Results – Total distance per match period



* indicates significant difference between backs and forwards, # indicated significant different from all other match periods. T, S, M, L and VL indicate effect sizes trivial (<0.2), small (0.2-0.5), medium (0.5-0.8), large (0.8-1.2) and very large (>1.2) respectively.




Results – High-intensity distance per match period



* indicates significant difference between backs and forwards, # indicates significant different from match period 2nd half Q4. T, S, M, L and VL indicate effect sizes trivial (<0.2), small (0.2-0.5), medium (0.5-0.8), large (0.8-1.2) and very large (>1.2) respectively.

Results – Match period effects sprint variables

Sprint and acceleration frequency are reduced in the 2nd half for forwards, but not for backs.



* indicates significant difference between backs and forwards. T, S, M, L and VL indicate effect sizes trivial (<0.2), small (0.2-0.5), medium (0.5-0.8), large (0.8-1.2) and very large (>1.2) respectively.


The effect of physical contact

↑ physical contact = ↓ total
↓ high-intensity running
distance (Johnston et al., 2014, Int J Sports Physiol
Perform, Epub)


Forwards experience ↑ contact
involvements than backs
(Deutsch et al., 2007, J Sport Sci 25:4)


Accelerometer data – backs
experience ↑ total (>5G) and
high-intensity (>8G) “impacts”
than forwards


Data doesn't fit fatigue model







Results – Effect of substitutes



Forward substitutes  sprint and acceleration frequency and high-intensity impacts


Back substitutes  sprint and acceleration frequency, but n = 3

Results – effect of forward substitutes



* indicates significant difference between whole game players and substitutes. T, S, M, L and VL indicate effect sizes trivial (<0.2), small (0.2-0.5), medium (0.5-0.8), large (0.8-1.2) and very large (>1.2) respectively.

Conclusions – running distance

Rugby union players  total (10%) and high-intensity (18%) running in 2nd half.

Similar results in soccer¹, rugby league² and rugby sevens³.

BUT

Rugby union work rates are much lower than other sports (~ 70 vs. $\sim 100 \text{ m} \cdot \text{min}^{-1}$)^{1,2,3}

1. Bradley and Noakes, 2013, J Sport Sci 31:15


2. Waldron et al., 2013, Int J Sports Physiol Perform 8:2

3. Higham et al., 2011, J Sci Med Sport 15




Conclusions – fatigue profile

Backs and forwards demonstrate differing fatigue profiles.




Pacing profile	
Forwards	Backs
“Slow positive”	“Flat”



Forwards progressively  total and high-intensity distance, maximum speed, sprint and acceleration frequency

Backs maintain total and high-intensity distance, maximum speed, sprint and acceleration frequency for majority of match

Conclusions – Impact of substitutes

substitute pacing strategy



Substitutes  match intensity by  high-intensity distance, acceleration frequency and high-intensity impacts.

Substitutes set a higher pacing strategy in the early part of their exercise bout

– a “**one bout, all out**” strategy

For the coach - Take home message

- Fatigue is evidenced by reductions in total and high intensity running distance and sprint and acceleration frequencies.
- Fatigue profile of forwards and backs is different
- Monitor high-intensity running distance to determine onset of fatigue
- Replacing fatigued players with substitutes is an effective method of maintaining playing intensity



Goodbye and thank you for listening!

Acknowledgements

Thank you to the players and staff of the
Golden Lions Rugby Union for their support
of the project

This research was partially funded by the
National Research Foundation.

Email: jasonctee@gmail.com

